

Mizoram Board
Class XI
Chemistry
Sample Paper – 1 Solution

Time: -3 hrs

Maximum Marks: - 70

1. This is because o – nitro phenol has intramolecular hydrogen bonding where as p- nitro phenol has intermolecular hydrogen bonding.
2. CO_2 has a larger bond angle than BF_3 . This is because CO_2 has a linear shape and the bond angle is 180° , BF_3 on the other hand has a trigonal planar geometry and hence the bond angle is 120° .
3. Energy change in climbing the hill is a state function. This is because it is independent of the path followed to reach the state.
4. $\text{NaH} \xrightarrow{\text{electrolysis}} \text{Na}^+ + \text{H}^-$
At anode: $2\text{H}^- - 2e^- \longrightarrow \text{H}_2(\text{g})$
5. Alkali metals have low ionization energies. They can lose electrons when light falls on them, and hence are used in photo electric cells.
6. The eclipsed conformation of propane is less stable and has more energy than the eclipsed conformation of ethane. This is because in propane there are additional interactions between C-H and C-C bond of methyl group.
7. $\text{O}_2\text{NCH}_2\text{CH}_2\text{O}^-$ will be more stable because $-\text{NO}_2$ group has electron withdrawing inductive effect or – I effect.
8. In Antarctica, ozone depletion is due to the formation of chlorine nitrate.
9.
 - (a) C will have the highest first ionization enthalpy since it has the smallest size and highest effective nuclear charge.
 - (b) C will have the most negative electron gain enthalpy because of its small size.
10. Statement a is correct and b is incorrect.
Statement b is incorrect because d sub shell can have a maximum of 10 electrons. Therefore it has 10 columns and not 8.

Or


(a) Since the element is in the third period $\therefore n=3$

The element is in the seventeenth group


\therefore The outermost configuration is: $3s^2 3p^5$

Therefore, the atomic number of the atom is 17.

(b) Cr (Z=24) has 5 electrons in the d sub shell. Its electronic configuration is:



11.



$$54 \text{ g of Al give H}_2 = 3 \times 22.4 \text{ L}$$

$$0.15 \text{ g of Al gives H}_2 = \frac{3 \times 22.4}{54} \times 0.15$$

$$= 0.186 \text{ L}$$

So, 0.186 L will be released at 1 bar pressure and 273 K. To calculate volume of H₂ at 20°C and 1 bar pressure,

$$V_1 = 0.186 \text{ L}$$

$$V_2 = ?$$


$$T_1 = 273 \text{ K}$$

$$T_2 = (20^\circ\text{C} + 273) \text{ K} = 293 \text{ K}$$

$$\frac{V_1}{T_1} = \frac{V_2}{T_2}$$

$$V_2 = \frac{0.186 \times 293}{273} = 0.1996 \text{ L} = 199.6 \text{ mL}$$

12. Ammonia will liquefy first because its critical temperature will be reached first. Liquefaction of CO₂ will require more cooling.





F₂ is the oxidizing agent and H₂O is reducing agent.

H₂O is getting oxidized to O₂ whereas F₂ is getting reduced to F⁻ ion.


14. A is Beryllium (Be)

B is Aluminum (Al)



15. Mg is present in chlorophyll.

N₂ reacts with Mg to form magnesium nitride.



Magnesium nitride


16. Molar mass of $\text{Na}_2\text{SO}_4 = (2 \times 23) + 32 + (4 \times 16) = 142 \text{ g mol}^{-1}$

$$\text{Mass \% of sodium} = \frac{2 \times 23}{142} \times 100 = 32.39\%$$


$$\text{Mass \% of sulphur} = \frac{32}{142} \times 100 = 22.53\%$$

$$\text{Mass \% of oxygen} = \frac{4 \times 16}{142} \times 100 = 45.07\%$$



17.



Therefore the structure of the compound will be -



18. Normally, rain water has a pH of about 5.6 due to the dissolution of CO_2 of the atmosphere into it.



When the pH of rain water falls below 5.6, it becomes acid rain.

$$\mathbf{19.} \quad x_{\text{C}_2\text{H}_5\text{OH}} = \frac{n_{\text{C}_2\text{H}_5\text{OH}}}{n_{\text{C}_2\text{H}_5\text{OH}} + n_{\text{H}_2\text{O}}}$$

For dilute solution, 1 L of solution can be nearly equal to 1 L of water.

$$n_{\text{H}_2\text{O}} = \frac{1000}{18} = 55.55 \text{ moles}$$

$$\frac{n_{\text{C}_2\text{H}_5\text{OH}}}{n_{\text{C}_2\text{H}_5\text{OH}} + 55.55} = 0.040$$

$$n_{\text{C}_2\text{H}_5\text{OH}} = 2.31 \text{ moles}$$

20.

(a) Atom is empty space. Being able to see the atom and structure of the atom itself means looking at empty space. Hence all of us will be seen as empty space. The wall will be seen as empty space. You and I will be seen as empty space. Hence, Kavita will not be able to see anything. In any other words she will be blind.

- (b) Praise to the almighty who has limited our abilities such that we are able to enjoy this colourful world.

21.

(a) 7

(b) This means that 6 electrons are present in p sub shell of the 4th shell

(c) $Z = 29$

Electronic configuration $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^1$

Or

(a) Number of electrons in 1 molecule of methane = $6 + 4 = 10$ electrons

Number of molecules in 1 mole of methane = 6.022×10^{23} molecules of methane

Number of electrons in 1 mole of methane = 6.022×10^{24} electrons

(b) $n = 3$

$l = 0 \text{ to } (n-1)$

= 0, 1, 2

For $l = 0,$

$m_l = 0$

For $l = 1$


$m_l = -1, 0, +1$


For $l = 2$

$m_l = -2, -1, 0, +1, +2$

22. The electronic configuration of S = $1s^2 2s^2 2p^6 3s^2 3p^4$

Sulphur undergoes sp^3d hybridisation.





23. (a)
$$K = \frac{[HI]^2}{[H_2]}$$

(b) According to Henderson's equation,

$$pOH = pK_b + \log \frac{[\text{salt}]}{[\text{base}]}$$

Also, $pK_b = -\log K_b$

$$= -\log 1.85 \times 10^{-5} = 4.733$$

$$pOH = 4.733 + \log \frac{0.2}{0.1}$$

$$= 4.733 + 0.3010 = 5.034$$

$$pH = 14 - pOH = 14 - 5.034 = 8.966$$



24.

(a) The equilibrium will shift the backward direction as the increase in temperature will be compensated by absorbing heat. It is an exothermic reaction.

(b) The equilibrium will shift in the forward direction since the reaction will shift to the direction of lesser number of moles.

(c) The equilibrium will shift in the forward direction so that additional SO_2 is used up.

25.



$$O = 15$$

$$O = 15$$

$$H = 5$$


$$H = 5$$

Oxygen and Hydrogen atoms are balanced.

26.

- (a) 3 - Methylpentanenitrile
- (b) 3-Chloropropanal
- (c) 4- Nitroaniline

27. (a) Order of stability





This is because $-\text{CH}_3$ group has electron releasing inductive effect or +I effect. Due to this, electron density increases on the negatively charged carbon and hence makes it more unstable. As the number of methyl groups increases the instability increases.

(b) The negatively charged carbon atom in a carbanion is sp^3 hybridised

28.



(a)

Compound (A) is $\text{H}_3\text{C}-\text{C}\equiv\text{C}-\text{C}_2\text{H}_5$



cis - isomer


(B)




trans - isomer

(C)



(b) Kolbe's electrolytic method -





Mechanism:



At anode:





At cathode:







At cathode hydrogen is liberated. At anode ethane is formed.

Or



↓ (i) O_3 (ii) $\text{Zn}/\text{H}_2\text{O}$





$$\Delta H = 177 \text{ kJ mol}^{-1}$$

$$\Delta S = 285 \text{ J K}^{-1} \text{ mol}^{-1} = 0.285 \text{ kJ K}^{-1} \text{ mol}^{-1}$$

$$T = 25^\circ\text{C} = 298 \text{ K}$$

$$\Delta G = \Delta H - T\Delta S$$





$$= 177 - (298 \times 0.285)$$

$$= +92.07 \text{ kJ mol}^{-1}$$


The reaction will be non-spontaneous.

This is because the value of ΔG is positive.


Or



Multiplying equation (2) by 6 and (3) by 3, and adding, (1 × 3)



Subtracting eqn (4) - eqn (1)




30.





- Boron has a very small size and has a very high sum of three ionisation enthalpies ($\text{IE}_1 + \text{IE}_2 + \text{IE}_3$). Therefore, it cannot lose its three electrons to form B^{3+} ions.
- AlBr_3 is predominantly a covalent compound. Even in molten state it does not have ions which can conduct electricity.
- B exhibits +3 oxidation state and can form stable BCl_3 . Thallium shows +3 oxidation state as well as +1 oxidation state but +1 oxidation state is more predominant than +3 oxidation state because of inert pair effect. Therefore, TiCl_3 is not stable. It can form stable TiCl .
- BCl_3 molecule has a symmetrical trigonal planar structure in which three B-Cl bonds are oriented at an angle of 120° to one another. The three bonds lie in one plane and the dipole moments of these bonds cancel one another giving net dipole moment zero.
- Electrical conductivity of Aluminium is twice as that of copper. On mass to mass basis, Al conducts electricity twice as Cu. Therefore, it is used in transmission cables.

Or

(a)



Dichlorodimethyl silicon



Boric acid